

Introduction to QTL mapping in model organisms

Karl W Broman

Department of Biostatistics
Johns Hopkins University


kbroman@jhsph.edu

www.biostat.jhsph.edu/~kbroman


Outline

- Experiments and data
- Goals, statistical structure
- Models
- ANOVA and interval mapping
- LOD thresholds
- CIs for QTL location
- Selection bias
- Multiple QTLs, epistasis
- The X chromosome
- Selective genotyping
- Covariates
- Non-normal traits
- The need for good data

Backcross experiment




Intercross experiment



Phenotype distributions

- Within each of the parental and F₁ strains, individuals are genetically identical.
- Environmental variation may or may not be constant with genotype.
- The backcross generation exhibits genetic as well as environmental variation.




Data

Phenotypes: y_i = trait value for individual i


Genotypes: x_{ij} = 0/1 if mouse i is BB/AB at marker j
(or 0/1/2, in an intercross)

Genetic map: Locations of markers

Genetic map




Genotype data



Goals

- Detect QTLs (and interactions between QTLs)
- Confidence intervals for QTL location
- Estimate QTL effects (effects of allelic substitution)

Statistical structure



The missing data problem:

Markers \longleftrightarrow QTL

The model selection problem:

QTL, covariates \longrightarrow phenotype


Models: Recombination

We assume no crossover interference.

⇒ Points of exchange (crossovers) are according to a Poisson process.

⇒ The $\{x_{ij}\}$ (marker genotypes) form a Markov chain

Example



Models: Genotype \longleftrightarrow Phenotype

Let y = phenotype
 g = whole genome genotype

Imagine a small number of QTLs with genotypes g_1, \dots, g_p .
(2^p distinct genotypes)

$$E(y|g) = \mu_{g_1, \dots, g_p} \quad \text{var}(y|g) = \sigma_{g_1, \dots, g_p}^2$$

Models: Genotype \longleftrightarrow Phenotype

Homoscedasticity (constant variance): $\sigma_g^2 \equiv \sigma^2$


Normally distributed residual variation: $y|g \sim N(\mu_g, \sigma^2)$.

Additivity: $\mu_{g_1, \dots, g_p} = \mu + \sum_{j=1}^p \Delta_j g_j$ ($g_j = 1$ or 0)

Epistasis: Any deviations from additivity.

The simplest method: ANOVA

- Also known as **marker regression**.
- Split mice into groups according to genotype at a marker.
- Do a t-test / ANOVA.
- Repeat for each marker.



Effect at a marker

Consider the case of a **single QTL** with effect $\Delta = \mu_{BB} - \mu_{AB}$.

Consider a marker linked to the QTL, with $r =$ recomb. frac.

Of individuals with marker genotype BB, mean phenotype is:

$$\mu_{BB}(1 - r) + \mu_{AB}r = \mu_{BB} - r\Delta$$

Of individuals with marker genotype AB, mean phenotype is:

$$\mu_{AB}(1 - r) + \mu_{BB}r = \mu_{AB} + r\Delta$$

Difference: $(\mu_{BB} - r\Delta) - (\mu_{AB} + r\Delta) = \Delta(1 - 2r)$

ANOVA at marker loci

Advantages

- Simple.
- Easily incorporates covariates.
- Easily extended to more complex models.
- Doesn't require a genetic map.


Disadvantages

- Must exclude individuals with missing genotype data.
- Imperfect information about QTL location.
- Suffers in low density scans.
- Only considers one QTL at a time.

Interval mapping (IM)

Lander & Botstein (1989)

- Take account of missing genotype data
- Interpolate between markers
- Maximum likelihood under a mixture model




Interval mapping (IM)

Lander & Botstein (1989)


- Assume a **single** QTL model.
- Each position in the genome, one at a time, is posited as the putative QTL.
- Let $z = 1/0$ if the (unobserved) QTL genotype is BB/AB.
Assume $y \sim N(\mu_z, \sigma)$
- Given genotypes at linked markers, $y \sim$ mixture of normal dist'ns with mixing proportion $\Pr(z = 1 | \text{marker data})$:

		QTL genotype	
M_1	M_2	BB	AB
BB	BB	$(1 - r_L)(1 - r_R)/(1 - r)$	$r_L r_R/(1 - r)$
BB	AB	$(1 - r_L)r_R/r$	$r_L(1 - r_R)/r$
AB	BB	$r_L(1 - r_R)/r$	$(1 - r_L)r_R/r$
AB	AB	$r_L r_R/(1 - r)$	$(1 - r_L)(1 - r_R)/(1 - r)$

The normal mixtures



- Two markers separated by 20 cM, with the QTL closer to the left marker.
- The figure at right show the distributions of the phenotype conditional on the genotypes at the two markers.
- The dashed curves correspond to the components of the mixtures.



Interval mapping (continued)

Let $p_i = \Pr(z_i = 1 | \text{marker data})$

$$y_i | z_i \sim N(\mu_{z_i}, \sigma^2)$$

$$\Pr(y_i | \text{marker data}, \mu_0, \mu_1, \sigma) = p_i f(y_i; \mu_1, \sigma) + (1 - p_i) f(y_i; \mu_0, \sigma)$$

$$\text{where } f(y; \mu, \sigma) = \exp[-(y - \mu)^2 / (2\sigma^2)] / \sqrt{2\pi\sigma^2}$$

Log likelihood: $l(\mu_0, \mu_1, \sigma) = \sum_i \log \Pr(y_i | \text{marker data}, \mu_0, \mu_1, \sigma)$

Maximum likelihood estimates (**MLEs**) of μ_0, μ_1, σ :

values for which $l(\mu_0, \mu_1, \sigma)$ is maximized.

EM algorithm

Dempster et al. (1977)

E step:

$$\begin{aligned} \text{Let } w^{(k+1)} &= \Pr(z_i = 1 | y_i, \text{marker data}, \hat{\mu}_0^{(k)}, \hat{\mu}_1^{(k)}, \hat{\sigma}^{(k)}) \\ &= \frac{p_i f(y_i; \hat{\mu}_1^{(k)}, \hat{\sigma}^{(k)})}{p_i f(y_i; \hat{\mu}_1^{(k)}, \hat{\sigma}^{(k)}) + (1 - p_i) f(y_i; \hat{\mu}_0^{(k)}, \hat{\sigma}^{(k)})} \end{aligned}$$

M step:

$$\begin{aligned} \text{Let } \hat{\mu}_1^{(k+1)} &= \sum_i y_i w_i^{(k+1)} / \sum_i w_i^{(k+1)} \\ \hat{\mu}_0^{(k+1)} &= \sum_i y_i (1 - w_i^{(k+1)}) / \sum_i (1 - w_i^{(k+1)}) \\ \hat{\sigma}^{(k+1)} &= [\text{not worth writing down}] \end{aligned}$$

The algorithm:

Start with $w_i^{(1)} = p_i$; iterate the E & M steps until convergence.

Example

Iteration	$\hat{\mu}_0$	$\hat{\mu}_1$	$\hat{\sigma}$	log likelihood
1	5.903	6.492	1.668	-770.752
2	5.835	6.562	1.654	-770.291
3	5.818	6.579	1.651	-770.264
4	5.815	6.583	1.650	-770.262
⋮	⋮	⋮	⋮	⋮
∞	5.813	6.584	1.649	-770.262

LOD scores



The LOD score is a measure of the **strength of evidence** for the presence of a QTL at a particular location.

$$\begin{aligned}\text{LOD}(z) &= \log_{10} \text{likelihood ratio comparing the hypothesis of a} \\ &\quad \text{QTL at position } z \text{ versus that of no QTL} \\ &= \log_{10} \left\{ \frac{\Pr(y|\text{QTL at } z, \hat{\mu}_{0z}, \hat{\mu}_{1z}, \hat{\sigma}_z)}{\Pr(y|\text{no QTL}, \hat{\mu}, \hat{\sigma})} \right\}\end{aligned}$$

$\hat{\mu}_{0z}, \hat{\mu}_{1z}, \hat{\sigma}_z$ are the MLEs, assuming a single QTL at position z .

No QTL model: The phenotypes are independent and identically distributed (iid) $N(\mu, \sigma^2)$.

An example LOD curve



Interval mapping

Advantages

- Takes proper account of missing data.
- Allows examination of positions between markers.
- Gives improved estimates of QTL effects.
- Provides pretty graphs.

Disadvantages

- Increased computation time.
- Requires specialized software.
- Difficult to generalize.
- Only considers one QTL at a time.

Haley-Knott regression

A quick approximation to Interval Mapping.


$$E(y \mid \text{QTL} = q) = \mu + \beta \mathbf{1}\{q = \text{AB}\}$$

$$E(y \mid \text{marker data}) = \mu + \beta \Pr(\text{QTL} = \text{AB} \mid \text{marker data})$$

- Regress y on $\Pr(\text{QTL} = \text{AB} \mid \text{marker data})$.
- **Pretend** that the residual variation is normally distributed.
- Calculate

$$\text{LOD}(\gamma) = (n/2) \log_{10} \left\{ \frac{\text{RSS}_0}{\text{RSS}_a(\gamma)} \right\}$$

Example



LOD thresholds

Large LOD scores indicate evidence for the presence of a QTL.

Q: How large is large?

→ We consider the distribution of the LOD score under the null hypothesis of no QTL.


Key point: We must make some adjustment for our examination of multiple putative QTL locations.

→ We seek the distribution of the *maximum* LOD score, genome-wide. The 95th %ile of this distribution serves as a **genome-wide LOD threshold**.


Estimating the threshold: simulations, analytical calculations, permutation (randomization) tests.

Null distribution of the LOD score

- Null distribution derived by computer simulation of backcross with genome of typical size.
- Solid curve: distribution of LOD score at any one point.
- Dashed curve: distribution of maximum LOD score, genome-wide.




Permutation tests




- Permute/shuffle the phenotypes; keep the genotype data intact.
- Calculate $\text{LOD}^*(z) \rightarrow M^* = \max_z \text{LOD}^*(z)$
- We wish to compare the observed M to the distribution of M^* .
- $\Pr(M^* \geq M)$ is a genome-wide P-value.
- The 95th %ile of M^* is a genome-wide LOD threshold.
- We can't look at all $n!$ possible permutations, but a random set of 1000 is feasible and provides reasonable estimates of P-values and thresholds.
- **Value:** conditions on observed phenotypes, marker density, and pattern of missing data; doesn't rely on normality assumptions or asymptotics.

Permutation distribution




1.5-LOD support interval



Selection bias

- The estimated effect of a QTL will vary somewhat from its true effect.
- Only when the estimated effect is large will the QTL be detected.
- Among those experiments in which the QTL is detected, the estimated QTL effect will be, on average, larger than its true effect.
- This is **selection bias**.
- Selection bias is largest in QTLs with small or moderate effects.
- The true effects of QTLs that we identify are likely smaller than was observed.



Implications of selection bias

- Estimated % variance explained by identified QTLs
- Repeating an experiment
- Congenics
- Marker-assisted selection

Multiple QTL methods

Why consider multiple QTLs at once?

- Reduce residual variation.
- Separate linked QTLs.
- Investigate interactions between QTLs (epistasis).

Issues:

- Missing genotype information
- The model selection problem

Data structure

y = phenotypes


m = observed marker genotypes

q = unobserved QTL genotypes

μ = model parameters

γ = QTL locations

H = QTL model



Likelihood vs. Bayes

$$l(H, \mu, \gamma | y, m) = \sum_q \Pr(y, q | m, \mu, \gamma, H)$$

Likelihood:

$$l(H | y, m) = \max_{\mu, \gamma} l(H, \mu, \gamma | y, m)$$

Bayes:

$$\Pr(H, \mu, \gamma | y, m) \propto l(H, \mu, \gamma | y, m) \Pr(\mu, \gamma | H) \Pr(H)$$


$$\Pr(H | y, m) = \iint \Pr(H, \mu, \gamma | y, m) d\mu d\gamma$$

Model selection


- **Select class of models**
 - Additive models
 - Add've plus pairwise interactions
 - Regression trees
- **Search model space**
 - Forward selection (FS)
 - Backward elimination (BE)
 - FS followed by BE
 - MCMC
- **Compare models**
 - Penalized likelihood (e.g., AIC, BIC)
 - Sequential permutation tests
 - Bayes (posterior probability)
- **Assess performance**
 - Maximize no. QTLs found; control false positive rate

Epistasis in a backcross

Additive QTLs




Interacting QTLs




Epistasis in an intercross

Additive QTLs



Interacting QTLs



Two-dimensional genome scan

Consider each pair of positions, (γ_1, γ_2)


Models

- Full
- Additive
- QTL 1
- QTL 2
- Null

Possible comparisons

- Full vs. null
- Full vs. additive
- Full vs. Best of QTL 1 & 2
- Add've vs. Best of QTL 1 & 2

Example



The X chromosome

In a backcross, the X chromosome may or may not be segregating.

$$(A \times B) \times A$$

Females: $X_{A \cdot B} X_A$

Males: $X_{A \cdot B} Y_A$

$$A \times (A \times B)$$

Females: $X_A X_A$

Males: $X_A Y_B$

The X chromosome

In an intercross, one must pay attention to the **paternal grandmother's genotype**.

$$(A \times B) \times (A \times B) \quad \text{or} \quad (B \times A) \times (A \times B)$$

Females: $X_{A \cdot B} X_A$

Males: $X_{A \cdot B} Y_B$


$$(A \times B) \times (B \times A) \quad \text{or} \quad (B \times A) \times (B \times A)$$

Females: $X_{A \cdot B} X_B$

Males: $X_{A \cdot B} Y_A$


Selective genotyping

- Save effort by only typing the most informative individuals (say, top & bottom 10%).
- Useful in context of a **single, inexpensive** trait.
- Tricky to estimate the effects of QTLs: use IM with **all** phenotypes.
- Can't get at interactions.
- Likely better to also genotype some random portion of the rest of the individuals.



Covariates

- **Examples:** treatment, sex, litter, lab, age.
- Control residual variation.
- Avoid confounding.
- Look for QTL \times environ't interactions
- Adjust before interval mapping (IM) versus adjust within IM.



Non-normal traits

- Standard interval mapping assumes normally distributed residual variation. (Thus the phenotype distribution is a mixture of normals.)
- **In reality:** we see dichotomous traits, counts, skewed distributions, outliers, and all sorts of odd things.
- Interval mapping, with LOD thresholds derived from permutation tests, generally performs just fine anyway.
- Alternatives to consider:
 - Nonparametric approaches (Kruglyak & Lander 1995)
 - Transformations (e.g., log, square root)
 - Specially-tailored models (e.g., a generalized linear model, the Cox proportional hazard model, and the model in Broman et al. 2000)

Check data integrity

The success of QTL mapping depends crucially on the integrity of the data.

- Segregation distortion
- Genetic maps / marker positions
- Genotyping errors (tight double crossovers)
- Phenotype distribution / outliers
- Residual analysis

Summary I

- **ANOVA** at marker loci (aka marker regression) is simple and easily extended to include covariates or accommodate complex models.
- **Interval mapping** improves on ANOVA by allowing inference of QTLs to positions between markers and taking proper account of missing genotype data.
- ANOVA and IM consider only single-QTL models. **Multiple QTL methods** allow the better separation of linked QTLs and are necessary for the investigation of epistasis.
- Statistical significance of LOD peaks requires consideration of the maximum LOD score, genome-wide, under the null hypothesis of no QTLs. **Permutation tests** are extremely useful for this.
- **1.5-LOD support intervals** indicate the plausible location of a QTL.
- Estimates of QTL effects are subject to **selection bias**. Such estimated effects are often too large.

Summary II

- The **X chromosome** must be dealt with specially, and can be tricky.
- **Study your data**. Look for errors in the genetic map, genotyping errors and phenotype outliers. But don't worry about them too much.
- **Selective genotyping** can save you time and money, but proceed with caution.
- **Study your data**. The consideration of covariates may reveal extremely interesting phenomena.
- Interval mapping works reasonably well even with **non-normal traits**. But consider transformations or specially-tailored models. If interval mapping software is not available for your preferred model, start with some version of ANOVA.

References

- Broman KW (2001) Review of statistical methods for QTL mapping in experimental crosses. *Lab Animal* 30:44–52
[A review for non-statisticians.](#)
- Doerge RW (2002) Mapping and analysis of quantitative trait loci in experimental populations. *Nat Rev Genet* 3:43–52
[A very recent review.](#)
- Doerge RW, Zeng Z-B, Weir BS (1997) Statistical issues in the search for genes affecting quantitative traits in experimental populations. *Statistical Science* 12:195–219
[Review paper.](#)
- Jansen RC (2001) Quantitative trait loci in inbred lines. In Balding DJ et al., *Handbook of statistical genetics*, John Wiley & Sons, New York, chapter 21
[Review in an expensive but rather comprehensive and likely useful book.](#)
- Lynch M, Walsh B (1998) *Genetics and analysis of quantitative traits*. Sinauer Associates, Sunderland, MA, chapter 15
[Chapter on QTL mapping.](#)
- Lander ES, Botstein D (1989) Mapping Mendelian factors underlying quantitative traits using RFLP linkage maps. *Genetics* 121:185–199
[The seminal paper.](#)

- Churchill GA, Doerge RW (1994) Empirical threshold values for quantitative trait mapping. *Genetics* 138:963–971
[LOD thresholds by permutation tests.](#)
- Kruglyak L, Lander ES (1995) A nonparametric approach for mapping quantitative trait loci. *Genetics* 139:1421–1428
[Non-parameteric interval mapping.](#)
- Broman KW (2003) Mapping quantitative trait loci in the case of a spike in the phenotype distribution. *Genetics* 163:1169–1175
[QTL mapping with a special model for a non-normal phenotype.](#)
- Miller AJ (2002) *Subset selection in regression*, 2nd edition. Chapman & Hall, New York.
[A good book on model selection in regression.](#)
- Strickberger MW (1985) *Genetics*, 3rd edition. Macmillan, New York, chapter 11.
[An old but excellent general genetics textbook with a very interesting discussion of epistasis.](#)