

How to display data badly

Using Microsoft Excel to
obscure your data and
annoy your readers

Inspiration

This lecture was inspired by

H Wainer (1984) How to display data badly. American Statistician
38(2):137-147

Dr. Wainer was the first to elucidate the principles of the bad display of data.

The now widespread use of Microsoft Excel has resulted in remarkable advances in the field.

General principles


The aim of good data graphics:

Display data accurately and clearly.

Some rules for displaying data badly:

- Display as little information as possible.
- Obscure what you do show (with chart junk).
- Use pseudo-3d and color gratuitously.
- Make a pie chart (preferably in color and 3d).
- Use a poorly chosen scale.
- Ignore sig figs.


Example 1



5


Example 2

Distribution of genotypes




6

Example 3




7

Example 4



8

Example 5




9

Example 6

N	$b/c = 10.0$		$b/c = 10.0$		$b/c = 100.0$	
	r^*	G	r^*	G	r^*	G
3	2	0.2	2	2.225	2	22.47499
4	2	0.26333	2	2.88833	2	29.13832
5	2	0.32333	3	3.54167	3	35.79166
6	3	0.38267	3	4.23767	3	42.78764
7	3	0.446	3	4.901	3	49.45097
8	3	0.50743	4	5.5765	4	56.33005
9	3	0.56743	4	6.26025	4	63.20129
10	4	0.62948	4	6.92358	4	69.86462

10

Example 7



11

Displaying data well

- Be accurate and clear.
- Let the data speak.
 - Show as much information as possible, taking care not to obscure the message.
- Science not sales.
 - Avoid unnecessary frills — esp. gratuitous 3d.
- In tables, every digit should be meaningful. Don't drop ending 0's.


12

Further reading

- ER Tufte (1983) The visual display of quantitative information. Graphics Press.
- ER Tufte (1990) Envisioning information. Graphics Press.
- ER Tufte (1997) Visual explanations. Graphics Press.
- WS Cleveland (1993) Visualizing data. Hobart Press.
- WS Cleveland (1994) The elements of graphing data. CRC Press.

Displaying distributions

Dotplots




- Few data points per group
- Possibly many groups


15

Histograms

Symmetric distribution




Skewed distribution



- Many data points per group
- Few groups
- Area of rectangle prop. to number of data points in interval
- I like **many** bins
(typically $2\sqrt{n}$)

16

Boxplots



- Many data points
- Possibly many groups
- Displays minimum, lower quartile, median, upper quartile and maximum

17

Summary statistics

Location/ Center

mean (i.e., average)
median
mode
geometric mean
harmonic mean

Scale

standard deviation (SD)
inter-quartile range (IQR)
range

Other

quantile / percentile
quartile
quintile

18

The means

$$\text{mean} = \frac{1}{n} \sum_{i=1}^n x_i = (x_1 + x_2 + \dots + x_n)/n$$

$$\text{geometric mean} = \sqrt[n]{\prod_{i=1}^n x_i} = \exp \left\{ \frac{1}{n} \sum_{i=1}^n \ln x_i \right\}$$


$$\text{harmonic mean} = 1 / \left\{ \frac{1}{n} \sum_{i=1}^n (1/x_i) \right\}$$

19


Measures of location/center

- Forget about the **mode**.
- The **mean** is **sensitive** to outliers.
(The balance point for the histogram.)
- The **median** is **resistant** to outliers.
- The **geometric mean** is used when a logarithmic transformation is appropriate (for example, when the distribution has a long right tail).
- The **harmonic mean** may be used when a reciprocal transformation is appropriate (very seldom).

20



21



22

A key point

The different possible measures of the “center” of the distribution are all **allowable**.

- Which is the best measure of the “typical” value (for your situation)?
- Be sure to make clear which “average” you use.

23


Standard deviation (SD)

$$\text{sample variance} = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2 = s^2$$


$$\begin{aligned} \text{sample SD} &= \sqrt{s^2} = s \\ &= \text{RMS (distance from average)} \\ &= \text{“typical” distance from the average} \\ &= \text{sort of like } \text{ave}\{|x_i - \bar{x}|\} \end{aligned}$$

$$\text{Note: } \bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$$

24





25



26

Logarithms


0.01	10^{-2}	1/4	2^{-2}
0.1	10^{-1}	1/2	2^{-1}
1	10^0	1	2^0
10	10^1	2	2^1
100	10^2	4	2^2
1000	10^3	8	2^3
10,000	10^4	16	2^4

log base 10 log base 2

30

Log transformation



31

mean = 100; SD = 30

